



## Factsheet 4: Management of Sexually Transmitted Infection (STI), as of Asar 2072 (July 2015)

### Background about STI in Nepal


- ✎ Key interventions for management of STI in Nepal are targeted Behavior Change Communication (BCC), condom promotion and distribution, diagnosis and treatment of STI (both syndromic and etiological management) and referral services.
- ✎ STI management services are available from Government Health Facilities and NGOs for key population.
- ✎ Nepal has been following WHO recommended approach for the management of STIs in patients with recognized signs and symptoms.
- ✎ The first National STI Case Management Guideline was developed in 1995 and the latest revision was made in 2014.
- ✎ Repeated Integrated Biological and Behavioral Surveillance (IBBS) Surveys are the main source of information for STI prevalence among key population in Nepal.

**Table 1: Self-reported STIs cases**

Regions	Female		Male	
	2006	2011	2006	2011
Eastern	0.2	0.9	0.0	0.0
Central	0.5	0.6	0.4	0.6
Western	0.1	0.1	0.0	0.3
Mid-Western	0.9	0.4	0.2	0.0
Far western	0.2	0.4	0.0	0.0

Source: NDHS 2006, 2011

**Figure 1: Annual Reported Cases of STI/STDs through HMIS reporting**



Source: Annual Report of DoHS

**For Further Information Please Contact:**


National Centre for AIDS and STD Control, Teku, Kathmandu

**Phone:** +977 - 1 - 4258219, 4261653, 4262753, **Email:** data@ncasc.gov.np

**Fax:** +977 - 1 - 4215 149 **Website:** www.ncasc.gov.np




**Figure 2: Prevalence of Active Syphilis among FSWs**



Source: Integrated Bio-Behavioural Surveillance (IBBS) Survey

**Figure 3: STI Prevalence among FSWs in 22 Terai Highway Districts**



Source: Integrated Bio-Behavioural Surveillance (IBBS) Survey

**For Further Information Please Contact:**


National Centre for AIDS and STD Control, Teku, Kathmandu

**Phone:** +977 - 1 - 4258219, 4261653, 4262753, **Email:** data@ncasc.gov.np

**Fax:** +977 - 1 - 4215 149 **Website:** www.ncasc.gov.np




Figure 4: Prevalence of Active Syphilis among MSM/TG in Kathmandu Valley



Source: Integrated Bio-Behavioural Surveillance (IBBS) Survey

Figure 4: STI Prevalence among MSM/TG in Kathmandu Valley



Source: Integrated Bio-Behavioural Surveillance (IBBS) Survey

**For Further Information Please Contact:**

National Centre for AIDS and STD Control, Teku, Kathmandu

**Phone:** +977 - 1 - 4258219, 4261653, 4262753, **Email:** data@ncasc.gov.np


**Fax:** +977 - 1 - 4215 149 **Website:** www.ncasc.gov.np



Government of Nepal  
Ministry of Health and Population  
**National Centre for AIDS and STD Control**  
Teku, Kathmandu




Figure 5: Prevalence of Active Syphilis among PWIDs in Kathmandu Valley



Source: Integrated Bio-Behavioural Surveillance (IBBS) Survey

Figure 6: Prevalence of Active Syphilis among PWIDs



Source: Integrated Bio-Behavioural Surveillance (IBBS) Survey

**For Further Information Please Contact:**

National Centre for AIDS and STD Control, Teku, Kathmandu

**Phone:** +977 - 1 - 4258219, 4261653, 4262753, **Email:** [data@ncasc.gov.np](mailto:data@ncasc.gov.np)

**Fax:** +977 - 1 - 4215 149 **Website:** [www.ncasc.gov.np](http://www.ncasc.gov.np)